

BEST AVAILABLE COPY

Serial No. 10/088,400

HANTKE et al.


0480/01219

APPENDIX I:

CLAIM AMENDMENTS:

Cancel Claims 5, 9 and 17 to 19, and amend Claims 1, 4, 11, 15, 16 and 20, as indicated in the following listing of the claims:

1. (currently amended) Rate-controlled release particles, comprising, in a polymer matrix consisting of a homo- or copolymer of N-vinylpyrrolidone, an active ingredient as a solid dispersion in the polymeric matrix consisting of a homo- or copolymer of N-vinylpyrrolidone and from 5 to 25% b.w. of hydroxypropyl methyl cellulose, and optionally further comprising a surfactant, and wherein the active ingredient is a compound of formula I



(I)

a N-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein

Y is CR⁵ or N;

A is CH, CR⁴ or N;

n is 0, 1, 2, 3 or 4;

Q is -NR¹R² or when Y is CR⁵ then Q may also be hydrogen;

R¹ and R² are each independently selected from hydrogen, hydroxy, C₁₋₁₂alkyl, C₁₋₁₂alkyloxy, C₁₋₁₂alkylcarbonyl, C₁₋₁₂alkyloxycarbonyl, aryl, amino, mono- or di(C₁₋₁₂alkyl)amino, mono- or di(C₁₋₁₂alkyl)aminocarbonyl

wherein each of the aforementioned C₁₋₁₂alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C₁₋₆alkyloxy, hydroxy-C₁₋₆alkyloxy, carboxyl, C₁₋₆alkyloxycarbonyl, cyano, amino, imido, aminocarbonyl, aminocarbonylamino, mono- or di(C₁₋₆alkyl)amino, aryl and Het; or

R¹ and R² taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C₁₋₁₂alkyl)aminoC₁₋₄-alkylidene;

R³ is hydrogen, aryl, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyl substituted with C₁₋₆alkyloxycarbonyl; and

051216

- 5 -

BEST AVAILABLE COPY

Serial No. 10/088,400

HANTKE et al.

0480/01219

each R^4 independently is hydroxy, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, or when Y is CR^5 then R^4 may also represent C_{1-6} alkyl substituted with cyano or amino carbonyl;

R^5 is hydrogen or C_{1-4} alkyl;

L is $-X^1-R^6$ or $-X^2-Alk-R^7$ wherein

R^6 and R^7 each independently are phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl, formyl, cyano, nitro, amino, and trifluoromethyl; or when Y is CR^5 then R^6 and R^7 may also be selected from phenyl substituted with one, two, three, four or five substituents each independently selected from aminocarbonyl, trihalomethyloxy and trihalomethyl; or when Y is N then R^6 and R^7 may also be selected from indanyl or indolyl, each of said indanyl or indolyl may be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl, formyl, cyano, nitro, amino, and trifluoromethyl;

X^1 and X^2 are each independently $-NR^3-$, $-NH-NH-$, $-N=N-$, $-O-$, $-S-$, $-S(=O)-$ or $-S(=O)_2-$;

Alk is C_{1-4} alkanediyl; or

when Y is CR^5 then L may also be selected from C_{1-10} alkyl, C_{3-10} alkenyl, C_{3-10} alkynyl, C_{3-7} cycloalkyl, or C_{1-10} alkyl substituted with one or two substituents independently selected from C_{3-7} cycloalkyl, indanyl, indolyl and phenyl, wherein said phenyl, indanyl and indolyl may be substituted with one, two, three, four or where possible five substituents each independently selected from halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, aminocarbonyl, C_{1-6} alkyloxycarbonyl, formyl, nitro, amino, trihalomethyl, trihalomethyloxy and C_{1-6} alkylcarbonyl;

aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, nitro and trifluoromethyl;

Het is an aliphatic or aromatic heterocyclic radical; said aliphatic heterocyclic radical is selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl

BEST AVAILABLE COPY

051216

- 6 -


Serial No. 10/088,400

HANTKE et al.

0480/01219

wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic heterocyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein each of said aromatic heterocyclic radical may optionally be substituted with hydroxy,





or a compound of formula II



(II)

the N-oxides, the pharmaceutically acceptable addition salts, quaternary amines and the stereochemically isomeric forms thereof, wherein

$-b^1=b^2-C(R^{2a})=b^3-b^4=$ represents a bivalent radical of formula



q is 0, 1, 2; or where possible q is 3 or 4;

R¹ is hydrogen, aryl, formyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyl substituted with formyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxycarbonyl;

R^{2a} is cyano, aminocarbonyl, mono- or di(methyl)aminocarbonyl, C₁₋₆alkyl substituted with cyano, aminocarbonyl or mono- or di(methyl)aminocarbonyl, C₂₋₆alkenyl substituted with cyano, or C₂₋₆alkynyl substituted with cyano;

each R² independently is hydroxy, halo, C₁₋₆alkyl optionally substituted with cyano or $-C(=O)R^6$, C₃₋₇cycloalkyl, C₂₋₆alkenyl optionally substituted with one or more halogen atoms or cyano, C₂₋₆alkynyl optionally substituted with one or more halogen atoms or cyano, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, $-S(=O)OR^6$,

051216

- 7 -

Serial No. 10/088,400

HANTKE et al.

0480/01219

$-\text{NH}-\text{S}(=\text{O})_p\text{R}^6$, $-\text{C}(=\text{O})\text{R}^6$, $-\text{NHC}(=\text{O})\text{H}$, $-\text{C}(=\text{O})\text{NHNH}_2$, $-\text{NEC}(=\text{O})\text{R}^6$,
 $-\text{C}(=\text{NH})\text{R}^6$ or a radical of formula



(c)

wherein each A independently is N, CH or CR^6 ;

B is NH, O, S or NR^6 ;

p is 1 or 2; and

R^6 is methyl, amino, mono- or dimethylamino or polyhalomethyl;

L is C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-7} cycloalkyl, whereby each of said aliphatic group may be substituted with one or two substituents independently selected from

* C_{3-7} cycloalkyl,

* indolyl or isoindolyl, each optionally substituted with one, two, three or four substituents each independently selected from halo, C_{1-6} alkyl, hydroxy, C_{1-6} alkyloxy, cyano, aminocarbonyl, nitro, amino, polyhalomethyl, polyhalomethyloxy and C_{1-6} alkylcarbonyl,

* phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R^2 ; or

L is $-\text{X}-\text{R}^3$ wherein

R^3 is phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R^2 ; and

X is $-\text{NH}^1-$, $-\text{NH}-\text{NH}-$, $-\text{N}=\text{N}-$, $-\text{O}-$, $-\text{C}(=\text{O})-$, $-\text{CHOH}-$, $-\text{S}-$, $-\text{S}(=\text{O})-$ or $-\text{S}(=\text{O})_2-$;

Q represents hydrogen, C_{1-6} alkyl, halo, polyhalo C_{1-6} alkyl or $-\text{NR}^4\text{R}^5$; and

R^4 and R^5 are each independently selected from hydrogen, hydroxy, C_{1-12} alkyl, C_{1-12} alkyloxy, C_{1-12} alkylcarbonyl, C_{1-12} alkyloxycarbonyl, aryl, amino, mono- or di(C_{1-12} alkyl)amino, mono- or di(C_{1-12} alkyl)aminocarbonyl

051216

- 8 -

Serial No. 10/088,400

HANTKE et al.

0480/01219

wherein each of the aforementioned C_{1-12} alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C_{1-6} alkyloxy, hydroxyc $_{1-6}$ alkyloxy, carboxyl, C_{1-6} alkyloxycarbonyl, cyano, amino, imino, mono- or di(C_{1-6} alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, $-S(O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(O)R^6$, $-C(=NH)R^6$, aryl and Het; or

R^4 and R^5 taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C_{1-12} alkyl)amino C_{1-4} -alkylidene;


Y represents hydroxy, halo, C_{3-7} cycloalkyl, C_{2-6} alkenyl optionally substituted with one or more halogen atoms, C_{2-6} alkynyl optionally substituted with one or more halogen atoms, C_{1-6} alkyl substituted with cyano or $-C(=O)R^6$, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C_{1-6} alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^6$, $-C(NH)R^6$ or aryl;

aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, cyano, nitro, polyhalo C_{1-6} alkyl and polyhalo C_{1-6} alkyloxy;

Het is an aliphatic or aromatic heterocyclic radical;

said aliphatic heterocyclic radical is selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic heterocyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein each of said aromatic heterocyclic radical may optionally be substituted with hydroxy,

or a compound of formula III



a N-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein

Serial No. 10/088,400

HANTKE et al.

0480/01219

$-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula

$-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$ (a-1);

$-\text{N}=\text{CH}-\text{CH}=\text{CH}-$ (a-2);

$-\text{N}=\text{CH}-\text{N}=\text{CH}-$ (a-3);

$-\text{N}=\text{CH}-\text{CH}=\text{N}-$ (a-4);

$-\text{N}=\text{N}-\text{CH}=\text{CH}-$ (a-5);

n is 0, 1, 2, 3 or 4; and in case $-a^1=a^2-a^3=a^4-$ is (a-1), then n may also be 5;

R^1 is hydrogen, aryl, formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkyl substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyl; and

each R^2 independently is hydroxy, halo, C_{1-6} alkyl optionally substituted with cyano or $-\text{C}(=\text{O})\text{R}^4$, C_{3-7} cycloalkyl, C_{2-6} alkenyl optionally substituted with one or more halogen atoms or cyano, C_{2-6} alkynyl optionally substituted with one or more halogen atoms or cyano, C_{1-6} alkyloxy, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C_{1-6} alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, $-\text{S}(=\text{O})_p\text{R}^4$, $-\text{NH}-\text{S}(=\text{O})_p\text{R}^4$, $-\text{C}(=\text{O})\text{R}^4$, $-\text{NHC}(=\text{O})\text{R}$, $-\text{C}(=\text{O})\text{NHNH}_2$, $\text{NHC}(=\text{O})\text{R}^4$, $-\text{C}(=\text{NH})\text{R}^4$ or a radical of formula



(c)

wherein each A independently is N, CH or CR^4 ;

B is NH, O, S or NR^4 ;

p is 1 or 2; and

R^4 is methyl, amino, mono- or dimethylamino or polyhalomethyl;

L is C_{4-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-7} cycloalkyl, whereby each of said aliphatic group may be substituted with one or two substituents independently selected from

* C_{3-7} cycloalkyl,

* indolyl or isoindolyl, each optionally substituted with one, two, three or four substituents each independently selected from halo, C_{1-6} alkyl, hydroxy, C_{1-6} alkyloxy, cyano, aminocarbonyl, nitro, amino, polyhalomethyl, polyhalomethyloxy and C_{1-6} alkylcarbonyl,

Serial No. 10/088,400

HANTKE et al.

0480/01219

* phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R²; or


L is -X-R³ wherein

R³ is phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with two, three, four or five substituents each independently selected from the substituents defined in R²; and

X is -NR¹-, -NH-NH-, -N=N-, -O-, -C(=O)-, -CHO-, -S-, -S(=O)- or -S(=O)₂-;

aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C₁₋₆alkyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, cyano, nitro, polyhaloC₁₋₆alkyl and polyhaloC₁₋₆alkyloxy,

or a compound of formula IV



the pharmaceutically acceptable acid addition salts and the stereochemically isomeric forms thereof, wherein

R¹ and R² are each independently selected from hydrogen; hydroxy; amino; C₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxy-carbonyl; Ar¹; mono- or di(C₁₋₆alkyl)amino; mono- or di(C₁₋₆alkyl)aminocarbonyl; dihydro-2(3H)-furanone; C₁₋₆alkyl substituted with one or two substituents each independently selected from amino, imino, aminocarbonyl, aminocarbonylamino, hydroxy, hydroxyc₁₋₆alkyloxy, carboxyl, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyloxy-carbonyl and thienyl; or

R¹ and R² taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C₁₋₆alkyl)aminoC₁₋₄-alkylidene;

R³ is hydrogen, Ar¹, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxy-carbonyl, C₁₋₆alkyl substituted with C₁₋₆alkyloxy-carbonyl; and

Serial No. 10/088,400

HANTKE et al.

0480/01219

R⁴, R⁵, R⁶, R⁷ and R⁸ are each independently selected from hydrogen, hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethyloxy;

L is C₁₋₁₀alkyl, C₃₋₁₀alkenyl; C₃₋₁₀alkynyl; C₃₋₇cycloalkyl, or

L is C₁₋₁₀alkyl substituted with one or two substituents independently selected from C₃₋₇cycloalkyl;

indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl;

phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl; and,

Ar¹ is phenyl, or phenyl substituted with one, two or three substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, nitro or trifluoromethyl;

with the proviso that compounds (a) to (o)

Co. No.	Alk	R ¹ /R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸
a	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	CH ₃	H	H	H	H
b	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	NO ₂	H	H
c	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	C ₆ H ₅	H	H	H	H	H
d	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	NO ₂	H	CH ₃	H	H
e	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	NH ₂	H	H
f	4-(2-methylpropyl)phenylmethyl	H/H	H	H	CF ₃	H	H	H
g	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	Cl	H	H
h	4-(2-methylpropyl)phenylmethyl	H/H	H	H	H	H	H	H
i	3,4-dimethoxyphenylmethyl	H/H	H	H	H	H	H	H
j	2,3-dimethoxyphenylmethyl	H/H	H	H	H	H	H	H
k	3,4-diethoxyphenylmethyl	H/H	H	H	H	H	H	H
l	2-(3,5-(1,1-dimethylethyl)-4-hydroxy-phenyl)ethyl	H/H	H	H	H	H	H	H
m	2-(3,5-(1,1-dimethylethyl)-4-hydroxy-phenyl)ethyl	H/H	H	H	t-Bu	OH	t-Bu	H
n	Phenylmethyl	H/H	H	CH ₃	H	H	H	H
o	Phenylmethyl	H/H	H	H	H	H	H	H

are not included,

or a compound of formula V


051216

- 12 -

Serial No. 10/088,400

HANTKE et al.

0480/01219



the *N*-oxide forms, the pharmaceutically acceptable acid addition salts and stereochemically isomeric forms thereof,

wherein

n is zero, 1, 2 or 3;


X is N or CH;

each *R*¹ independently is halo, nitro, cyano, amino, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy or trifluoromethyl;

*R*² is hydrogen; C₃₋₇alkenyl; C₃₋₇alkynyl, aryl; C₃₋₇cycloalkyl; C₁₋₆alkyl or C₁₋₆alkyl substituted with hydroxy, C₁₋₄alkyloxy, C₃₋₇cycloalkyl or aryl;

*R*³ and *R*⁴ each independently are hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl or aryl; or


*R*³ and *R*⁴ taken together form a bivalent radical -*R*³-*R*⁴- of formula:



wherein *R*^{5a}, *R*^{5b}, *R*^{5c}, *R*^{5d} each independently are hydrogen, C₁₋₆alkyl or aryl; and

aryl is phenyl or phenyl substituted with one, two or three substituents selected from halo, nitro, cyano, amino, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy or trifluoromethyl,

or a compound of formula VI



the *N*-oxides, the stereochemically isomeric forms thereof, and the pharmaceutically acceptable acid addition salts, wherein A and B taken together form a bivalent radical of formula:

Serial No. 10/088,400

HANTKE et al.

0480/01219

- N=CH- (a),
- CH=N- (b),
- CH₂-CH₂- (c),
- CH=CH- (d),
- C(=O)-CH₂- (e),
- CH₂-C(=O)- (f),

in the bivalent radicals- of formula (a) and (b) the hydrogen atom may be replaced by C₁₋₆alkyl; in the bivalent radicals of formula (c), (d), (e), (f), one or two hydrogen atoms may be replaced by C₁₋₆alkyl;

R¹ is hydrogen, C₁₋₆alkyl or halo;

R² is hydrogen or halo;

R³ is hydrogen; C₁₋₈alkyl; C₃₋₆cycloalkyl; or C₁₋₈alkyl substituted with hydroxy, oxo, C₃₋₆cycloalkyl or aryl;

Het is a heterocycle selected from the group consisting of pyridine; pyridine substituted with one or two substituents selected from C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, trihalomethyl, amino, mono- or di(C₁₋₆alkyl)amino or aryl;

pyrimidine; pyrimidine substituted with one or two substituents selected from C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, trihalomethyl, amino, mono- or di(C₁₋₆alkyl)-amino or aryl;

tetrazole; tetrazole substituted with C₁₋₆alkyl or aryl;

triazole; triazole substituted with one or two substituents selected from C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, trihalomethyl, amino, mono- or di(C₁₋₆alkyl)-amino;

thiadiazole; thiadiazole substituted with one or two substituents selected from C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, trihalomethyl, amino, mono- or di(C₁₋₆alkyl)-amino;

oxadiazole substituted with one or two substituents selected from C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, trihalomethyl, amino, mono- or di(C₁₋₆alkyl)amino;

imidazole; imidazole substituted with one or two substituents selected from C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, trihalomethyl, amino, mono- or di(C₁₋₆alkyl)amino;

thiazole; thiazole substituted with one or two substituents selected from C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, trihalomethyl, amino, mono- or di(C₁₋₆alkyl)amino;

Serial No. 10/088,400

HANTKE et al.

0480/01219

oxazole; oxazole substituted with one or two substituents selected from C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, trihalomethyl, amino, mono- or di(C₁₋₆alkyl)amino;

aryl is phenyl or phenyl substituted with C₁₋₆alkyl or halo, and the heterocyclic radical "Het" is bound to the sulfur atom via a carbon atom.

2. (original) Particles according to claim 1, wherein the copolymer of N-vinylpyrrolidone is a copolymer with vinyl acetate.
3. (canceled)
4. (currently amended) Particles according to claim 1, which comprise a surfactant and wherein the surfactant is a PEG-n-hydrogenated castor oil, or a low molecular weight polyoxyethylene polyoxypropylene block copolymer.
5. (canceled)
6. (previously presented) Particles according to claim 1, further comprising citric acid in amounts of up to 5% b.w.
7. (previously presented) Particles according to claim 1, wherein the homo- or copolymer of N-vinylpyrrolidone is used in amounts of from 40 to 70% b.w. of the total weight of the dosage form.
8. (original) Particles according to claim 7, wherein the homo- or copolymer of N-vinylpyrrolidone is used in amounts of from 50 to 65 % b.w..
9. (canceled)
10. (previously presented) Particles according to claim 1, wherein the controlled release is a sustained release.
11. (currently amended) Particles according to claim 10, further comprising the hydroxypropyl methyl cellulose in amounts of from 5 to 10 % b.w..
12. (previously presented) Particles according to claim 1, obtained by forming a homogeneous mixture of the components in the form of a melt, extruding said mixture and shaping of the extrudate.
13. (previously presented) Particles according to claim 1, comprising a compound selected from the group consisting of

Serial No. 10/088,400

HANTKE et al.

0480/01219

4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidyl]amino]benzonitrile;

4-[[2-[(cyanophenyl)amino]-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;

4-[[4-amino-5-chloro-6-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;

4-[[5-chloro-4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;

4-[[5-bromo-4-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidin]amino]benzonitrile;

4-[[4-amino-5-chloro-6-[(4-cyano-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;

4-[[5-bromo-6-[(4-cyano-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;

4-[[4-amino-5-chloro-6-[(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;

4-[[4-amino-5-bromo-6-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;

4-[[4-[(2,4,6-trimethylphenyl)amino]-1,3,5-triazin-2-yl]amino]benzonitrile;

4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-1,3,5-triazin-2-yl]amino]benzonitrile;

4-[[4-[(2,6-dichlorophenyl)methyl]-6-(hydroxyamino)-1,3,5-triazin-2-yl]amino]benzonitrile;

1[4-[[4-[[4-[(2,4-difluorophenyl)-4-(1H-1,2,4-triazol-1-yl-methyl)-1,3-dioxolan-2-yl]methoxy]phenyl]-1-piperazinyl]-phenyl]-3-(1-methylethyl)-2-imidazolidinone;

(-)-[2S-[2alpha, 4alpha(S*)]]-4-[[4-[[4-[[2-(4-chlorophenyl)-2-[[[(4-methyl-4H-1,2,4-triazol-3-yl)thio]methyl]-1,3-dioxolan-4-yl]methoxyl]phenyl]-1-piperazinyl]phenyl]-2,4-dihydro-2-(1-methyl-propyl)-3H-1,2,4-triazol-3-one,

a N-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof.

14. (previously presented) Pharmaceutical dosage form, comprising particles according to a claim 1.
15. (currently amended) Pharmaceutical dosage forms according to claim ~~13~~ 14, further comprising one or more pharmaceutically acceptable excipients.

051216

- 16 -

Serial No. 10/088,400

HANTKE et al.

0480/01219

16. (currently amended) Particles according to claim 1 ~~4~~, which meet one or both of the following requirements:
- the surfactant has a HLB-value of from 10 to 18;
 - the surfactant is present in the particles in an amount of from 5 to 20% by weight.
17. - 19. (canceled)
20. (currently amended) Particles according to claim 1, consisting essentially of the active ingredient,
from 40 to 70% by weight of the a homo- or copolymer of N-vinylpyrrolidone,
from 5 to 20% by weight of the surfactant,
up to 5% by weight of citric acid, and
~~up from 5~~ to 25% by weight of hydroxypropyl methyl cellulose.
21. (previously presented) Particles according to claim 20, wherein the surfactant has a HLB-value of from 10 to 18.
22. (previously presented) Particles according to claim 21, wherein the surfactant is a PEG-n-hydrogenated castor oil and/or a low molecular weight polyoxyethylene polyoxypropylene block copolymer.
23. (previously presented) Particles according to claim 1, obtained by a process comprising forming a homogeneous mixture of the components in the form of a melt, extruding said melt and shaping the obtained extrudate.
24. (previously presented) Particles according to claim 16, obtained by a process comprising forming a homogeneous mixture of the components in the form of a melt, extruding said melt and shaping the obtained extrudate.
25. (previously presented) Particles according to claim 20, obtained by a process comprising forming a homogeneous mixture of the components in the form of a melt, extruding said melt and shaping the obtained extrudate.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.